

Bufo Regularis lucn

~~Ranger's toad—Wikipedia Sclerophrys regularis (African common toad) | BioLib.cz Amietophrynus regularis—Wikipedia tiếng Việt Sclerophrys kisolensis—Wikipedia Ranger's toad—Wikipedia (PDF) Phylogeography of the African Common Toad ... Sclerophrys poweri—Wikipedia Bufo regularis—Wikimedia Commons AmphibiaWeb—Sclerophrys regularis Bufo—Wikipedia Pantterikonna—Wikipedia Bufo Regularis lucn Amietophrynus regularis—Wikipedia Common toad—Wikipedia Species Page of Sclerophrys mauritanica—Amphibians ... Sclerophrys regularis—Wikispecies Sclerophrys kerinyagae—Wikipedia Sclerophrys regularis—Wikipedia~~

Ranger's toad - Wikipedia

Frost et al., 2006 agrupan tentativamente a la especie dentro del nuevo género Amietophrynus, donde la especie tipo es Bufo regularis Reuss, 1833; y cuya etimología hace honor Jean-Louis Amiet, un influyente herpetólogo del África occidental y phrynos, del griego, sapo.

Sclerophrys regularis (African common toad) | BioLib.cz

This page was last edited on 24 January 2015, at 15:34. Files are available under licenses specified on their description page. All structured data from the file and property namespaces is available under the Creative Commons CC0 License; all unstructured text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply.

Amietophrynus regularis - Wikipedia tiếng Việt

Phylogeography of the African Common Toad, Amietophrynus regularis , Based on Mitochondrial DNA Sequences: Inferences Regarding the Cape Verde Population and Biogeographical Patterns

Bookmark File PDF Bufo Regularis Iucn

Sclerophrys kisoensis - Wikipedia

Sclerophrys poweri is a species of toad in the family Bufonidae. It is found in Angola, Botswana, Namibia, and South Africa. Its natural habitats are dry savanna, moist savanna, subtropical or tropical dry shrubland, subtropical or tropical dry lowland grassland, swamps, freshwater marshes, intermittent freshwater marshes, arable land, pastureland, and ponds.

Ranger's toad - Wikipedia

Description Edit. True toads have in common stocky figures and short legs, which make them relatively poor jumpers. Their dry skin is thick and warty. Behind their eyes, Bufo species have wart-like structures, the parotoid glands. These glands distinguish the true toads from all other tailless amphibians.

(PDF) Phylogeography of the African Common Toad ...

An *Amietophrynus regularis* in uska species han Anura nga syahan ginhulagway ni Reuss hadton 1833. An *Amietophrynus regularis* in nahilalakip ha genus nga *Amietophrynus*, ngan familia nga Bufonidae. Ginklasipika han IUCN an species komo diri gud kababarak-an. Waray hini subspecies nga nakalista. Mga kasarigan

Sclerophrys poweri - Wikipedia

Bufo neglectus Rochebrune, 1884 *Bufo regularis regularis* — Loveridge, 1936 *Amietophrynus regularis* — Frost et al. , 2006 [comb. nov. Ohler & Dubois, 2016]

Bufo regularis - Wikimedia Commons

Pantterikonna (*Sclerophrys regularis*, *Amietophrynus regularis* eli *Bufo regularis*) on Afrikassa elävä rupikonnalaji. Se kasvaa 6–13 cm mittaiseksi, ja sen iho on rupikonnille tyypillisen nystermäpintainen. Sitä tavataan Saharan eteläpuolisessa Afrikassa, savannilla, metsänreunoissa ja

Bookmark File PDF Bufo Regularis lucn

viljelysmailla, usein jokien läheisyydessä.. Pantterikonna pidetään myös lemmikkeinä terraarioissa.

AmphibiaWeb - Sclerophrys regularis

Bufo kisolensis Loveridge, 1932 Amietophrynus kisolensis (Loveridge, 1932) Sclerophrys kisolensis occurs in mature, undisturbed montane forests at elevations of 1,500–3,000 m (4,900–9,800 ft) above sea level , possibly wider.

Bufo - Wikipedia

Hewitt considered the Raucous Toad, along with most other large toads of Southern Africa, as subspecies of the widespread African Toad Bufo regularis (a species which is now considered absent from Southern Africa, predominantly occurring in subsaharan Africa north of the equator).

Pantterikonna - Wikipedia

Amietophrynus regularis Reuss, 1833 ... Bufo regularis. Amietophrynus regularis là một loài cóc trong họ Bufonidae. Chúng được tìm thấy ở Angola, Bénin, Burkina Faso, ... Amietophrynus regularis 2006 IUCN Red List of Threatened Species. Truy cập ngày 19 tháng 1 năm 2011.

Bufo Regularis lucn

Bufo regularis Reuss, 1833 Amietophrynus regularis (Reuss, 1833) Sclerophrys regularis , commonly known as the African common toad , square-marked toad , African toad , Egyptian toad , African bouncing toad (due to the bouncing motion) and Reuss's toad , is a species of toad in the family Bufonidae .

Amietophrynus regularis - Wikipedia

Bufo regularis Reuss, 1833. Espesye sa baki nga una nga gihulagway ni Reuss ni adtong 1833 ang

Bookmark File PDF Bufo Regularis Iucn

Amietophrynus regularis. Ang *Amietophrynus regularis* sakop sa kahenera nga *Amietophrynus* sa kabanay nga *Bufo*idae. Giklaseklase sa IUCN ang espesye sa kinaminosang kalabotan. Pagka karon wala pay siak nga nalista ubos niini niya. ...

Common toad - Wikipedia

In 1964, Poynton raised *Bufo rangeri* of Hewitt, 1935 to full species status, distinguishing it from co-occurring related forms. More recently, Frost and co-authors (2006) assigned this species to the new genus *Amietophrynus* , as part of a global revision of amphibian taxonomy based largely on DNA sequence data.

Species Page of Sclerophrys mauritanica - Amphibians ...

Scientific synonyms. *Bufo regularis* Reuss, 1833 *Amietophrynus regularis* (Reuss, 1833) *Bufo regularis regularis* *Bufo amoenus* Rochebrune, 1884 *Bufo buccinator* Rochebrune, 1884 *Bufo guineensis* Günther, 1859 *Rana mosaica* Seetzen, 1855 *Bufo neglectus* Rochebrune, 1884 *Bufo nubicus* Fitzinger, 1826 *Bufo sauvagei* Rochebrune, 1884

Sclerophrys regularis - Wikispecies

Amietophrynus kerinyagae. Least Concern (IUCN 3.1) *Amietophrynus kerinyagae* is a species of toad in the family *Bufo*idae . It is found in Ethiopia, Kenya, Tanzania, and Uganda . Its natural habitats are subtropical or tropical moist montane forests, subtropical or tropical high-altitude shrubland, subtropical or tropical high-altitude...

Sclerophrys kerinyagae - Wikipedia

The common toad, European toad, or in Anglophone parts of Europe, simply the toad (*Bufo bufo*, from Latin *bufo* "toad"), is an amphibian found throughout most of Europe (with the exception of Ireland, Iceland, and some Mediterranean islands), in the western part of North Asia, and in a small

Bookmark File PDF Bufo Regularis lucn

portion of Northwest Africa.

Sclerophrys regularis - Wikipedia

According to Inger & Greenberg (1956), male *B. regularis* have most often a single, sometimes paired, lateral slit in the floor of the mouth that forms the connection to the median, subgular vocal sac.

Copyright code : d129fc04fbdf440549baea5696eeb249.